

What is laser peripheral iridotomy?

An iridotomy is a small hole created with the laser beam in the outer part of the iris (coloured part of the eye).

A laser peripheral iridotomy is performed to treat or prevent a sudden rise in pressure inside the eye (acute glaucoma). A sustained rise in pressure can cause a permanent loss of side and central vision. The iridotomy allows fluid to circulate freely within the eye minimising the risk of a sight-threatening pressure rise.

There are three reasons for this procedure:

- To prevent an attack of acute glaucoma.
- To treat an attack of acute glaucoma.
- To treat chronic glaucoma where the drainage channel is very narrow.

This laser procedure is performed to preserve sight. It will not restore or improve the sight.

Risks

Complications are uncommon. The most common complications are a rise in pressure and some inflammation shortly after the treatment. The pressure will be checked before you go home and if very high, you will need extra treatment in the form of drops, tablets or in some cases an injection,

You will also need to remain in the eye clinic until the pressure has dropped to a safe level.

Other risks include:

- Bleeding in front of the iris;
- Clouding of the cornea (window) of the eye;
- Cataract.

The day of your treatment

The treatment takes approximately 15 minutes, but please allow between one and two hours at the hospital. Your treatment will be carried out in the eye clinic; on your arrival the doctor will explain the procedure, ask you to sign a consent form and numb the eye with drops. The procedure is performed with the patient sitting and resting their chin on an instrument similar to that used to examine the eye. A contact lens is placed on the eye to steady it and focus the laser beam. You will see some bright flashes of light, hear a clicking noise and may feel slight discomfort.

After your treatment

The doctor will instil some more eye drops and the pressure will be checked one hour after the treatment. You may then return home.

Your eye may become pink and sore and the vision is often disturbed for the rest of the day. For this reason patients are strongly advised not to drive home. Apart from driving, you can resume normal activities immediately. You will be prescribed additional medication for a few days to prevent inflammation and a rise in pressure. **You should continue to use all your anti-glaucoma medication unless specifically instructed otherwise.**

An appointment will be made for a check-up in the eye clinic shortly afterwards. A number of patients will require more than one laser procedure.




Moorfields Eye Clinic
Bedford Hospital NHS Trust
Kempston Road
Bedford
MK42 9DJ
01234 792 643

Bedford Hospital **NHS**
NHS Trust



Ophthalmology

YAG Laser Peripheral Iridotomy Treatment of Glaucoma.



Information for patients

Date of publication: January 2010
Date for review: January 2012
© Bedford Hospital NHS Trust